

The background of the slide is a close-up photograph of red poppies and their green seed pods. The flowers are in various stages of bloom, with some fully open and others as buds. The green pods are prominent, showing their characteristic ribbed texture. The overall color palette is dominated by the red of the petals and the green of the pods, set against a blurred background of more foliage.

MANAGEMENT OF OPIOID USE DISORDER: Medication Assisted Treatment

*C. Luke Peterson, DO
Clinical Assistant Professor*

*Hillary Tamar, MD
Family Medicine Resident*

*University of Arizona
College of Medicine – Phoenix*


Disclosure

- I hereby declare that the content for this activity, including any presentation on therapeutic options, is well balanced, unbiased, and to the extent evidenced based.
 - My partner/spouse and I have no financial relationships with commercial entities producing, marketing, re-selling, or distributing health care goods or services consumed by, or used on, patients relevant to the content I am planning, developing, presenting, or evaluating.
-

Current Toll and Trends

3


- Overdose in 2017
 - 70,237 drug overdose
 - 59% involved synthetic opioids
 - Arizona with largest relative rate increase in synthetic opioid overdose (122%)
 - 1.8/100,000 deaths (2016)
 - 4/100,000 deaths (2017)
- Substances cut with fentanyl:
 - Street oxycodone, street benzodiazepines, heroin, cocaine, methamphetamine



Current Toll and Trends

4

- Overdose in 2017
 - 70,237 drug overdose
 - 59% involved synthetic opioids
 - Arizona with largest relative rate increase in synthetic opioid overdose (122%)
 - 1.8/100,000 deaths (2016)
 - 4/100,000 deaths (2017)
- Substances cut with fentanyl:
 - Street oxycodone, street benzodiazepines, heroin, cocaine, methamphetamine



Objectives

1

Describe medication
assisted treatment (MAT)

Addiction •
Sociopsychobiologic



2

Discuss MAT efficacy, length
of treatment, pregnancy, and
common perceptions



3

Discuss access to MAT
including telemedicine



DSM-V Criteria – Opioid Use Disorder and the 3C's

Loss of Control	Using larger amounts over a longer period than initially intended
	Persistent desire or inability to cut down or control opioid use
	Spending a lot of time to obtain, use, or recover from use
Craving	Cravings or strong desire to use
Use Despite Consequences	Failure to fulfill obligations at work, school, or home due to use
	Continued use despite persistent or recurrent social or interpersonal problems related
	Activities are given up or reduced because of use
	Recurrent use in situations that are physically hazardous
	Continue use despite physical or psychological problems related to opioids
Physiologic Dependence	Tolerance*
	Withdrawal*

Mild 2-3, Moderate 4-5, Severe ≥ 6


***Does not count toward Use Disorder if prescribed and taken as directed**

Why do we care about models?

- What causes [blank]?
- What are the treatment and goals?
- Who is responsible for treatment outcomes?
- How does the health system drive care?

BIOMEDICAL

The U.S. pays most per capita for health care and ranks one of the lowest: Life expectancy vs. health expenditure, 1970 → 2015.



BIOMEDICAL

BIO

PSYCHO

SOCIAL

BIOMEDICAL

BIO

PSYCHO

SOCIAL





Sociopsychobiologic approach to substance use disorder

13



Food, Water, Shelter/Housing
Financial, Insurance, Transportation
Domestic Violence
Human Trafficking

Family/Peer Support
Prevention
Destigmatizing Campaign
Criminal Justice Reform

Screening and Treat Mental Health Disorder
Counseling and Behavioral Support
Coping Skills / Resilience
Relapse Prevention

MAT: Methadone or Buprenorphine
Depo-naltrexone
Primary Care / Preventative Medicine

Multimodal Pain Mgmt
Harm Reduction

Medication Assisted Treatment

14

- Goals of treatment:
 - Regain control
 - Decrease opioid withdrawal
 - Decrease cravings
 - Decrease use opioids
 - Block subsequent use of opioids
 - Improve quality of life of individual and community
- Length of treatment
 - Individualized - "As long as is helpful" - no artificial limits
 - SAMHSA – at least 12 months



Medication Assisted Treatment

15

- FDA Approved Medications
 - Methadone
 - Buprenorphine
 - Suboxone, Subutex, Zubsolv, Probuphine, Sublocade
 - Depo-naltrexone (Vivitrol)
-



Socioeconomic Impacts of MAT

- Mortality
 - MAT lowers risk of death while in treatment, specifically death from opioid and other drug overdoses, trauma, and suicide
 - Not on MAT leads to:
 - 2.5 times risk of dying from any cause
 - 8 times risk of overdose death
 - In a study of 3789 patients, there were 113 overdose deaths
 - 61 before treatment
 - 24 during treatment
 - 28 after treatment
 - Save 25 lives if you treat 1,000 with methadone over 1 year



Socioeconomic Impacts of MAT

17

- Treatment impact
 - MAT **increases retention** in treatment
 - Methadone > buprenorphine
 - MAT **decreases drug use** and positive UDS



Socioeconomic Impacts of MAT

18

- Crime

- Involvement in crime and the amount of crime committed during periods of addiction is dramatically higher than during periods of non-addiction
- During MAT, rates of criminal convictions drop to less than half
 - Acquisitive, drug selling, and violent crimes
- Patients in continuous treatment have the fewest convictions



Socioeconomic Impacts of MAT

- DUI

- In one study, 78% of those who tested positive for heroin had been arrested previously for drunken or drugged driving
- During MAT, convictions for DUI were reduced by 40%
- Total convictions for men decreased by 35%, for women, 60%



Socioeconomic Impacts of MAT

20

- Economics

- Both methadone and buprenorphine are more cost-effective than no drug therapy in dependent opioid users
 - Methadone - £12,584 /QALY (£12584 = \$16,347)
 - Buprenorphine - £30,048/QALY (£30,048 = \$39,034)
- Annual commercial health care cost for patients on methadone is 50% less than non-methadone
 - Methadone - \$7,163
 - Non-methadone substance treatment - \$14,157
 - No substance treatment - \$18,694



Socioeconomic Impacts of MAT

21

- HIV Transmission
 - Between 5-10% of all HIV infections are due to injection drug use
 - With MAT, there is a **54% reduction in HIV transmission risk**
 - Multifactorial
 - Less unprotected sex
 - Less sexual partners
 - Less injection drug use



- Depo-Naltrexone
 - Given 7-14 days after last opioid
 - Harder to initiate (30% of patients drop out)
 - Mortality rate lower in treatment, not as favorable as methadone or buprenorphine during and after treatment
 - Retention non-inferior to buprenorphine at 3 mo
 - Treatment retention needs more research



Patient Vignette



Medication Assisted Treatment

- Pregnancy
 - American College of Obstetrics and Gynecology
 - MAT is **gold standard treatment**
 - Neonatal abstinence may occur and is treatable
 - 60% stopped MAT by 6 months postpartum
 - More developmental impact if parent using while child is growing up
 - **Goal is long term chronic disease management**



Detox or MAT?

	Detoxification	MAT
Treatment Retention	10-40%	70-95%
Opioid Positive UDS	50-80%	20-50%


- Flexible **high dose methadone** (60-100 mg) or **high dose buprenorphine** (≥ 16 mg) most effective at retention and preventing return to use
- *Can you be in recovery on MAT?*



Access to Medication Assisted Treatment

26

- Methadone maintenance (1974)
- Drug Abuse Treatment Act (2000)
 - 48% of waived physicians were prescribing buprenorphine to five patients or fewer
- Lack of MAT access in US
 - OUD rate 891/100,000
 - Methadone capacity 119/100,000
 - Buprenorphine capacity 420/100,000
 - 43% of counties have no DATA physicians
 - 2% of physicians had waiver and 90% were in urban areas
- AZ - 40% of treatment centers offer MAT




Access to Medication Assisted Treatment

27

- Methadone maintenance (1974)
- Drug Abuse Treatment Act (2000)
 - 48% of waived physicians were prescribing buprenorphine to five patients or fewer
- Lack of MAT access in US
 - OUD rate 891/100,000
 - Methadone capacity 119/100,000
 - Buprenorphine capacity 420/100,000
 - 43% of counties have no DATA physicians
 - 2% of physicians had waiver and 90% were in urban areas
- AZ - 40% of treatment centers offer MAT

Figure 1. US counties with physicians with waivers to prescribe buprenorphine.



Note: data source: Drug Enforcement Administration, July 2012. Map date: September 2013.

Access to Medication Assisted Treatment

28

- MAT in UME/GME/CME
 - ADHS Pain and Addiction
- MAT in Primary Care
 - Sociopsychobiologic support
- MAT in substance use treatment centers
- MAT offered to persons in criminal justice
- Telemedicine
 - More likely to engage in counseling
 - Increased retention in treatment



Objectives

1

Describe MAT

Addiction •
Sociopsychobiologic



2

MAT decreases mortality,
drug use, and crime.

MAT in pregnancy!

No duration limits



3

AZ still needs to increase
access to MAT



Works Cited

- Bell, James, et al. "Comparing Retention in Treatment and Mortality in People after Initial Entry to Methadone and Buprenorphine Treatment." *Addiction*, vol. 104, no. 7, 2009, pp. 1193-1200., doi:10.1111/j.1360-0443.2009.02627.x.
- Bukten, Anne, et al. "Driving under the Influence (DUI) among Patients in Opioid Maintenance Treatment (OMT): a Registry-Based National Cohort Study." *Addiction*, vol. 108, no. 11, 2013, pp. 1954-1961., doi:10.1111/add.12275.
- Bukten, Anne, et al. "Engagement with Opioid Maintenance Treatment and Reductions in Crime: a Longitudinal National Cohort Study." *Addiction*, vol. 107, no. 2, 2011, pp. 393-399., doi:10.1111/j.1360-0443.2011.03637.x.
- Center for Disease Control and Prevention. Opioid Data Analysis and Resources. <https://www.cdc.gov/drugoverdose/data/analysis.html>. Accessed on March 9, 2019.
- CDC/NCHS, National Vital Statistics System, Mortality. CDC WONDER, Atlanta, GA: US Department of Health and Human Services, CDC; 2018. <https://wonder.cdc.gov/>.
- Committee on Obstetric Practice. Committee Opinion No. 711: Opioid Use and Opioid Use Disorder in Pregnancy. *Obstet Gynecol*. 2017 Aug;130(2):e81-e94. doi:10.1097/AOG.0000000000002235. PubMed PMID: 28742676.
- Clausen, Thomas, et al. "Mortality Prior to, during and after Opioid Maintenance Treatment (OMT): A National Prospective Cross-Registry Study." *Drug and Alcohol Dependence*, vol. 94, no. 1-3, 2008, pp. 151-157., doi:10.1016/j.drugalcdep.2007.11.003.
- Connock, M, et al. "Methadone and Buprenorphine for the Management of Opioid Dependence: a Systematic Review and Economic Evaluation." *Health Technology Assessment*, vol. 11, no. 9, 2007.
- Dunn KE, Sigmon SC, Strain EC, Heil SH, Higgins ST. The association between outpatient buprenorphine detoxification duration and clinical treatment outcomes: a review. *Drug Alcohol Depend*. 2011 Dec 1;119(1-2):1-9. doi: 10.1016/j.drugalcdep.2011.05.033. Epub 2011 Jul 8. Review. PubMed PMID: 21741781; PubMed Central PMCID: PMC3205338. Degenhardt, Louisa, et al. "Mortality among Clients of a State-Wide Opioid Pharmacotherapy Program over 20 Years: Risk Factors and Lives Saved." *Drug and Alcohol Dependence*, vol. 105, no. 1-2, 2009, pp. 9-15., doi:10.1016/j.drugalcdep.2009.05.021.

Works Cited

- Gossop, Michael, et al. "Reductions in Criminal Convictions after Addiction Treatment: 5-Year Follow-Up." *Drug and Alcohol Dependence*, vol. 79, no. 3, 2005, pp. 295-302., doi:10.1016/j.drugalcdep.2005.01.023.
- Jones CM, Campopiano M, Baldwin G, McCance-Katz E. National and State Treatment Need and Capacity for Opioid Agonist Medication-Assisted Treatment. *Am J Public Health*. 2015 Aug;105(8):e55-63. doi: 10.2105/AJPH.2015.302664. Epub 2015 Jun 11. PubMed PMID: 26066931; PubMed Central PMCID: PMC4504312.
- Ma J, Bao YP, Wang RJ, Su MF, Liu MX, Li JQ, Degenhardt L, Farrell M, Blow FC, Ilgen M, Shi J, Lu L. Effects of medication-assisted treatment on mortality among opioids users: a systematic review and meta-analysis. *Mol Psychiatry*. 2018 Jun 22. doi: 10.1038/s41380-018-0094-5. [Epub ahead of print] PubMed PMID: 29934549.
- Mattick RP, Breen C, Kimber J, Davoli M. Buprenorphine maintenance versus placebo or methadone maintenance for opioid dependence. *Cochrane Database Syst Rev*. 2014 Feb 6;(2):CD002207. doi: 10.1002/14651858.CD002207.pub4. Review. PubMed PMID: 24500948.
- McCarty D, Perrin NA, Green CA, Polen MR, Leo MC, Lynch F. Methadone maintenance and the cost and utilization of health care among individuals dependent on opioids in a commercial health plan. *Drug Alcohol Depend*. 2010 Oct 1;111(3):235-40. doi: 10.1016/j.drugalcdep.2010.04.018. PubMed PMID: 20627427; PubMed Central PMCID: PMC2950212.
- "Methadone Reduces Risk of HIV Transmission in People Who Inject Drugs, Say Experts." *The BMJ*, 4 Oct. 2012, www.bmj.com/press-releases/2012/10/04/methadone-reduces-risk-hiv-transmission-people-who-inject-drugs-say-expert.
- Pain and Addiction Curriculum. Arizona Department of Health. 2018, <https://www.azdhs.gov/documents/audiences/clinicians/curriculum/arizona-pain-addiction-curriculum-august.pdf>
- Rosenblatt RA, Andrilla CH, Catlin M, Larson EH. Geographic and specialty distribution of US physicians trained to treat opioid use disorder. *Ann Fam Med*. 2015 Jan-Feb;13(1):23-6. doi: 10.1370/afm.1735. PubMed PMID: 25583888; PubMed Central PMCID: PMC4291261.
- Tesema L, Marshall J, Hathaway R, Pham C, Clarke C, Bergeron G, Yeh J, Soliman M, McCormick D. Training in office-based opioid treatment with buprenorphine in US residency programs: A national survey of residency program directors. *Subst Abus*. 2018;39(4):434-440. doi: 10.1080/08897077.2018.1449047. Epub 2018 May 14. PubMed PMID: 29513136.